

Rituxan (Rituximab) Patient Referral and Prescription Sheet Return Signed Rx via Fax to: 888.837.2716

Date:	To:		From:				
Phone:		Fax:	Numbe	ber of Pages:			
			Patient Information				
Patient Name:				DOB:		Height:	
Allergies:						Weight	•
Medication Order							
Rituximab:			Wedleadon Oraci				
\square Rituximab: 375mg x BSA (m ²) = IV every for courses.							
Titte Airing X	55, ((/	cvery	101 600	. 505.		
☐ Rituximab:							Refills
-First Rituximab IV dose	is to be administer	ed in a control	led environment setting th	nen subsequent doses in	the home set	ting	
-Dilute Rituxan to a final	concentration of 1	mg/mL - 4 mg	g/mL with either 0.9% Sod	ium Chloride, or 5% Dext	rose		
Diagnosis ICD-10		Diagnosis		ICD-10			
☐ Rheumatoid Arthritis M06		M06. 9	☐ Wegener's Granulomatosis			M31.3	
☐ Chronic Lymphocytic Leukemia (CLL)		C91.10	☐ Pemphigus Vulgaris (PV)			L10.0	
■ Non-Hodgkin's Lymp	homa (NHL)	C85. 9	Other:				
☐ Neuromyelitis Optica	[Devic]	G36.0					
IV Access Device: Peripheral Central Lab order: CBC with diff CMP							
Pre-medications:			If applicable, flush IV access device per KabaFusion protocol:				
Pre-medications:			ii applicable, ilusii iv ac	cess device per Kabaru	Sion protoco	I.	
☐ Diphenhydramine:			Access	NS	sion protoco	Heparir	n
☐ Diphenhydramine: ☐ Acetaminophen:	□ PO □ IV		• • •		10 U/mL 1-2	Heparir	
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone:	□ PO □ IV □ IV		Access	NS		Heparir 2 mL after l	ast NS flush
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone: ☐ Other pre-meds:		– b Dituuissab	Access Peripheral	NS 1-3 mL before/after use	10 U/mL 1-2	Heparir 2 mL after l 5 mL after la	ast NS flush ast NS flush
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone: ☐ Other pre-meds: ☐ Hydration: Infuse 500 mL		– h Rituximab	Access Peripheral Midline, Central (non-port),	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use.	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5	Heparir 2 mL after l 5 mL after la he blood d mL after la	ast NS flush ast NS flush Iraw st NS flush
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone: ☐ Other pre-meds: ☐ Hydration: Infuse 500 mL infusion	PO IV IV of Normal Saline with		Access Peripheral Midline, Central (non-port), PICC Implanted Port	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use. 20mL after blood draw	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5 5 mL after t	Heparir 2 mL after l 5 mL after la he blood d mL after la he blood d	ast NS flush ast NS flush lraw st NS flush Iraw
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone: ☐ Other pre-meds: ☐ Hydration: Infuse 500 mL infusion ☐ Anaphylaxis Protocol: Ep	PO IV IV of Normal Saline with		Access Peripheral Midline, Central (non-port), PICC	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use. 20mL after blood draw 5-10 mL before/after use	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5 5 mL after t 10 U/mL 3-5	Heparir 2 mL after l 5 mL after la he blood d mL after la he blood d 5 mL after la	ast NS flush ast NS flush Iraw st NS flush Iraw ast NS flush
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone: ☐ Other pre-meds: ☐ Hydration: Infuse 500 mL infusion ☐ Anaphylaxis Protocol: Ep Adult: 0.3 mg Childr Administer epinephrine IN	of Normal Saline with sinephrine Auto-Injection: 0.15 mg M in the event of ana	ector dual pack: phylaxis.	Access Peripheral Midline, Central (non-port), PICC Implanted Port Tunneled	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use. 20mL after blood draw 5-10 mL before/after use 20mL after blood draw	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5 5 mL after t 10 U/mL 3-5 5 mL after t	Heparir 2 mL after l 5 mL after la he blood d mL after la he blood d 5 mL after la	ast NS flush ast NS flush lraw st NS flush lraw ast NS flush
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone: ☐ Other pre-meds: ☐ Hydration: Infuse 500 mL infusion ☐ Anaphylaxis Protocol: Ep Adult: 0.3 mg Childr	of Normal Saline with sinephrine Auto-Injection: 0.15 mg M in the event of ana	ector dual pack: phylaxis.	Access Peripheral Midline, Central (non-port), PICC Implanted Port	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use. 20mL after blood draw 5-10 mL before/after use 20mL after blood draw 5-10 mL before/after use	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5 5 mL after t 10 U/mL 3-5	Heparir 2 mL after l 5 mL after la he blood d mL after la he blood d 5 mL after la	ast NS flush ast NS flush Iraw st NS flush Iraw ast NS flush
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone: ☐ Other pre-meds: ☐ Hydration: Infuse 500 mL infusion ☐ Anaphylaxis Protocol: Ep Adult: 0.3 mg Childr Administer epinephrine IN May repeat x 1 as needed	of Normal Saline with of Normal Saline with the Point of Saline with the Point of Saline with the event of Saline with th	ector dual pack: phylaxis. _I r	Access Peripheral Midline, Central (non-port), PICC Implanted Port Tunneled Groshong PICC, Midline	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use. 20mL after blood draw 5-10 mL before/after use 20mL after blood draw 5-10 mL before/after use 10 mL after blood draw	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5 5 mL after t 10 U/mL 3-5 5 mL after t None	Heparir 2 mL after I 5 mL after Ia he blood d mL after Ia he blood d 5 mL after Ia he blood d	ast NS flush ast NS flush raw st NS flush Iraw ast NS flush Iraw
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone: ☐ Other pre-meds: ☐ Hydration: Infuse 500 mL infusion ☐ Anaphylaxis Protocol: Ep Adult: 0.3 mg Childr Administer epinephrine IN May repeat x 1 as needed ☐ authorize KabaFusion and	of Normal Saline with Normal Saline Auto-Injecten: 0.15 mg M in the event of ana Normal Saline	phylaxis. o act as an agent	Access Peripheral Midline, Central (non-port), PICC Implanted Port Tunneled	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use. 20mL after blood draw 5-10 mL before/after use 20mL after blood draw 5-10 mL before/after use 10 mL after blood draw insurance prior authorization	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5 5 mL after t 10 U/mL 3-5 5 mL after t None	Heparin 2 mL after la 5 mL after la he blood d mL after la he blood d 5 mL after la he blood d	ast NS flush ast NS flush draw st NS flush draw ast NS flush draw
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone: ☐ Other pre-meds: ☐ Hydration: Infuse 500 mL infusion ☐ Anaphylaxis Protocol: Ep Adult: 0.3 mg Childr Administer epinephrine IN May repeat x 1 as needed I authorize KabaFusion and future fills of the same presc	of Normal Saline with Normal Saline Auto-Injecten: 0.15 mg M in the event of ana Normal Saline	phylaxis. o act as an agent	Access Peripheral Midline, Central (non-port), PICC Implanted Port Tunneled Groshong PICC, Midline and initiate and execute any	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use. 20mL after blood draw 5-10 mL before/after use 20mL after blood draw 5-10 mL before/after use 10 mL after blood draw insurance prior authorizations designation at any time by	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5 5 mL after t 10 U/mL 3-5 5 mL after t None	Heparin 2 mL after la 5 mL after la he blood d mL after la he blood d 5 mL after la he blood d	ast NS flush ast NS flush draw st NS flush draw ast NS flush draw
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone: ☐ Other pre-meds: ☐ Hydration: Infuse 500 mL infusion ☐ Anaphylaxis Protocol: Ep Adult: 0.3 mg Childr Administer epinephrine IN May repeat x 1 as needed ☐ authorize KabaFusion and	of Normal Saline with Normal Saline Auto-Injecten: 0.15 mg M in the event of ana Normal Saline	phylaxis. o act as an agent	Access Peripheral Midline, Central (non-port), PICC Implanted Port Tunneled Groshong PICC, Midline and initiate and execute any	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use. 20mL after blood draw 5-10 mL before/after use 20mL after blood draw 5-10 mL before/after use 10 mL after blood draw insurance prior authorization	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5 5 mL after t 10 U/mL 3-5 5 mL after t None	Heparin 2 mL after la 5 mL after la he blood d mL after la he blood d 5 mL after la he blood d	ast NS flush ast NS flush draw st NS flush draw ast NS flush draw
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone: ☐ Other pre-meds: ☐ Hydration: Infuse 500 mL infusion ☐ Anaphylaxis Protocol: Ep Adult: 0.3 mg Childr Administer epinephrine IN May repeat x 1 as needed ☐ authorize KabaFusion and future fills of the same presc Prescriber Signature: ☐	of Normal Saline with Normal Saline Auto-Injecten: 0.15 mg M in the event of ana Normal Saline	phylaxis. o act as an agent	Access Peripheral Midline, Central (non-port), PICC Implanted Port Tunneled Groshong PICC, Midline and initiate and execute any	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use. 20mL after blood draw 5-10 mL before/after use 20mL after blood draw 5-10 mL before/after use 10 mL after blood draw insurance prior authorization is designation at any time by	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5 5 mL after t 10 U/mL 3-5 5 mL after t None	Heparin 2 mL after la 5 mL after la he blood d mL after la he blood d 5 mL after la he blood d	ast NS flush ast NS flush draw st NS flush draw ast NS flush draw
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone: ☐ Other pre-meds: ☐ Hydration: Infuse 500 mL infusion ☐ Anaphylaxis Protocol: Ep Adult: 0.3 mg Childr Administer epinephrine IN May repeat x 1 as needed I authorize KabaFusion and future fills of the same presc	of Normal Saline with Normal Saline Auto-Injecten: 0.15 mg M in the event of ana Normal Saline	phylaxis. o act as an agent	Access Peripheral Midline, Central (non-port), PICC Implanted Port Tunneled Groshong PICC, Midline and initiate and execute any	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use. 20mL after blood draw 5-10 mL before/after use 20mL after blood draw 5-10 mL before/after use 10 mL after blood draw insurance prior authorizations designation at any time by	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5 5 mL after t 10 U/mL 3-5 5 mL after t None	Heparin 2 mL after la 5 mL after la he blood d mL after la he blood d 5 mL after la he blood d	ast NS flush ast NS flush draw st NS flush draw ast NS flush draw
☐ Diphenhydramine: Acetaminophen: Methylprednisolone: Other pre-meds: Hydration: Infuse 500 mL infusion ☐ Anaphylaxis Protocol: Ep Adult: 0.3 mg Childr Administer epinephrine IN May repeat x 1 as needed I authorize KabaFusion and future fills of the same presc Prescriber Signature: Print Prescriber Name:	of Normal Saline with IV	phylaxis. o act as an agent t listed above. I u	Access Peripheral Midline, Central (non-port), PICC Implanted Port Tunneled Groshong PICC, Midline and initiate and execute any nderstand that I can revoke th	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use. 20mL after blood draw 5-10 mL before/after use 20mL after blood draw 5-10 mL before/after use 10 mL after blood draw insurance prior authorizations designation at any time by	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5 5 mL after t 10 U/mL 3-5 5 mL after t None on process for tile providing writt	Heparin 2 mL after la 5 mL after la he blood d mL after la he blood d 5 mL after la he blood d	ast NS flush ast NS flush draw st NS flush draw ast NS flush draw
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone: ☐ Other pre-meds: ☐ Hydration: Infuse 500 mL infusion ☐ Anaphylaxis Protocol: Ep Adult: 0.3 mg	of Normal Saline with IV	phylaxis. o act as an agent t listed above. I u	Access Peripheral Midline, Central (non-port), PICC Implanted Port Tunneled Groshong PICC, Midline and initiate and execute any nderstand that I can revoke th	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use. 20mL after blood draw 5-10 mL before/after use 20mL after blood draw 5-10 mL before/after use 10 mL after blood draw insurance prior authorizations designation at any time by	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5 5 mL after t 10 U/mL 3-5 5 mL after t None on process for till providing writt	Heparin 2 mL after la 5 mL after la he blood d mL after la he blood d 5 mL after la he blood d	ast NS flush ast NS flush draw st NS flush draw ast NS flush draw
☐ Diphenhydramine: ☐ Acetaminophen: ☐ Methylprednisolone: ☐ Other pre-meds: ☐ Hydration: Infuse 500 mL infusion ☐ Anaphylaxis Protocol: Ep Adult: 0.3 mg Childr Administer epinephrine IN May repeat x 1 as needed ☐ authorize KabaFusion and future fills of the same presc ☐ Prescriber Signature: ☐ Print Prescriber Name: ☐ Rx Order – include dos ☐ Patient Demographics	of Normal Saline with IV	phylaxis. property act as an agent tilisted above. I under the property act and the property action, frequency, information. We	Access Peripheral Midline, Central (non-port), PICC Implanted Port Tunneled Groshong PICC, Midline and initiate and execute any nderstand that I can revoke th	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use. 20mL after blood draw 5-10 mL before/after use 20mL after blood draw 5-10 mL before/after use 10 mL after blood draw insurance prior authorizations designation at any time by	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5 5 mL after t 10 U/mL 3-5 5 mL after t None on process for till providing writt	Heparin 2 mL after la 5 mL after la he blood d mL after la he blood d 5 mL after la he blood d	ast NS flush ast NS flush draw st NS flush draw ast NS flush draw
□ Diphenhydramine: □ Acetaminophen: □ Methylprednisolone: □ Other pre-meds: □ Hydration: Infuse 500 mL infusion □ Anaphylaxis Protocol: Ep Adult: 0.3 mg Childr Administer epinephrine IN May repeat x 1 as needed □ I authorize KabaFusion and future fills of the same presc Prescriber Signature: Print Prescriber Name: □ Rx Order – include dos □ Patient Demographics □ H & P OR progress no	of Normal Saline with IV	phylaxis. phylaxis. p act as an agent t listed above. I u DOCUMENT ation, frequency, information. We nosis, clinical sta	Access Peripheral Midline, Central (non-port), PICC Implanted Port Tunneled Groshong PICC, Midline and initiate and execute any nderstand that I can revoke th	NS 1-3 mL before/after use 5-10 ml before/after use 5-20 ml after blood draw 5-10 mL before/after use. 20mL after blood draw 5-10 mL before/after use 20mL after blood draw 5-10 mL before/after use 10 mL after blood draw insurance prior authorizations designation at any time by	10 U/mL 1-2 10 U/mL 3-5 5 mL after t 100 U/mL 5 5 mL after t 10 U/mL 3-5 5 mL after t None on process for till providing writt	Heparin 2 mL after la 5 mL after la he blood d mL after la he blood d 5 mL after la he blood d	ast NS flush ast NS flush draw st NS flush draw ast NS flush draw

CONFIDENTIALITY NOTICE

The following includes confidential, proprietary information that is the sole exclusive property of KabaFusion Holdings, LLC. No rights in, relating to, or derived from such information are assigned or otherwise transferred by this document, and the recipient of such information is subject to obligations of secrecy to and for the benefit of KabaFusion Holdings, LLC. Any unauthorized use or disclosure of such information is strictly prohibited. This message, together with any attachments, is intended only for the use of the individual or entity to which it is addressed and may contain information that is confidential and prohibited from disclosure. If you are not the intended recipient, you are hereby notified that any dissemination, or copying of this message, or any attachment, is strictly prohibited. If you have received this message in error, please notify the original sender immediately by telephone or by return fax and shred this document along with any other documents. Thank you.